

Relativistic and QED energy shifts in positronium ion

M. Grigorescu and G. W. F. Drake

Department of Physics
University of Windsor
Windsor, Ontario, Canada

Abstract: The leading relativistic and QED corrections to the ground state energy of the three-body system $e^-e^+e^-$ are calculated numerically using a Hylleraas correlated basis set. The accuracy of the nonrelativistic variational ground state wave function is discussed with respect to the convergence properties at the increase of the basis dimension and to the variance of the energy expectation value. Recent progress in the numerical procedure used to calculate expectation values for products of various physical operators is presented. It is shown that the nonrelativistic ground state energy can be calculated with an accuracy below the level width. The corrections to this energy include the lowest order Breit interaction, the vacuum polarization potential, one and two photon exchange contributions, the annihilation interaction, and spin-spin contact terms. The relativistic effects and the residual interactions considered here decrease the one electron binding energy from the nonrelativistic value of 0.012 005 070 232 980 10(3) a.u. to 0.011 981 051 246(2) a.u..

PACS: 45.50.Jf,36.10.Dr,31.15.-p,12.38.Bx,02.70.-c

I. Introduction

The positronium negative ion (Ps^-) is the simplest system composed of three equal mass fermions, $e^-e^+e^-$, bounded only by electromagnetic interactions. Similar examples of three-body systems, bounded by increasingly complex interactions, are provided by three-quark systems such as the proton and neutron, and three-nucleon systems, as the ^3H , ^3He nuclei.

In a nonrelativistic approach, accurate numerical approximations to the bound eigenstates of three quantum particles interacting by Coulomb forces can be obtained by using the Rayleigh-Ritz variational method. A suitable set of coordinates and basis states for the three-body problem was proposed by Hylleraas [1] during the early days of quantum mechanics, and it was used to calculate the ground state energy of the helium atom. With respect to this set, the matrix elements of various two-body operators can be expressed in analytical form [2], and extensive high-precision calculations become feasible [3] [4].

The relativistic quantum many-body problem can be approached either from the field theory, or by using a Schrödinger equation with an "action at a distance" type Hamiltonian, defined by quantizing the classical relativistic system [5]. Though, a puzzling result in classical mechanics is the no interaction theorem [6], which apparently rules out any instantaneous action-at-a-distance Hamiltonian. This theorem states that in a classical many-body system, the relativistic invariance of the equations of motion (the physical laws) is compatible with the "manifest relativistic invariance", of the world lines determined by these equations, only if there is no interaction between the particles. However, this strong result can be avoided if the interacting particles have a structure, as the condition of manifest invariance becomes ambiguous [7] [8].

The approach to the bound state problem based on field theory leads to a relativistically invariant Bethe-Salpeter equation [9], p. 196. In the case of two relativistic electrons, approximate Lorentz invariance to the first order is introduced by the Breit interaction, which can be seen as the quantum correspondent to the Darwin term in classical electromagnetism [10].

In the helium atom, the two electrons move in the Coulomb field created by a composite, heavy nucleus, which to a first approximation can be considered as center of mass (CM). The case of Ps^- is different, because all three particles are elementary, have the same mass, and move to the same degree

with respect to the CM.

The existence of a bound ground state in the $e^-e^+e^-$ system was predicted by Wheeler [11] and was observed by Mills [12] passing a positron beam through a thin carbon film in vacuum. The measured $\text{Ps}^- \rightarrow (2\gamma)e^-$ decay rate $\lambda = 2.09(9) \text{ nsec}^{-1}$ [12] corresponds to a Ps^- lifetime of 0.478 nsec, intermediate between that of para (singlet) Ps (0.125 nsec) and ortho (triplet) Ps (140 nsec) [13].

Accurate nonrelativistic numerical calculations for the ground-state properties of Ps^- are presented in refs. [14] to [17]. The autoionization states have been studied in [18], while several low-lying resonances have been predicted recently [19], by using a combination between the stochastic variational method (SVM) with correlated Gaussians and the complex scaling method.

The accuracy of the Ps^- groundstate wave functions given by SVM in a Gaussian basis, was studied by comparison to the direct solution of the Schrödinger equation in [16]. It was shown that despite the fact that in SVM the convergence properties of the expectation values for most operators are better, the wave function is less accurate.

In this work, the accuracy of the Ps^- nonrelativistic variational ground state is studied by using beside the convergence properties of the energy with basis size, also the variance of the Hamiltonian. The numerical procedure used to calculate matrix elements is presented in Sect. II. It is shown that in agreement with [16], the variance is larger than the accuracy resulting from convergence. Estimates of the relativistic correction terms and the leading QED corrections are presented in Sect. III. Tables containing the expectation values of singular operators appearing in the correction terms, such as p^4 and delta functions are given in the Appendix. The main results and the concluding remarks are summarized in Sect. IV.

II. The Nonrelativistic Quantum Three-Body Problem

The nonrelativistic Hamiltonian of the three-body system $e^-e^+e^-$ (or $e^+e^-e^+$) is

$$H_0 = \left(-\frac{1}{2}\nabla_1^2 - \frac{1}{2}\nabla_2^2 - f\nabla_1 \cdot \nabla_2 - \frac{1}{r_1} - \frac{1}{r_2} + \frac{1}{r_{12}} \right) f \text{ a.u.} \quad (1)$$

where $f = \mu/m$, $\mu = m/2$ is the reduced mass, $\nabla_i \equiv \partial_{\vec{r}_i}$, $\vec{r}_i = \vec{R}_i/a_\mu$,

$i = 1, 2, 3$ denote the position vectors in the CM frame of the two electrons ($i = 1, 2$), and of the positron ($i = 3$) in a_μ units, while $r = |\vec{r}_1 - \vec{r}_2|$, $r_1 = |\vec{r}_1 - \vec{r}_3|$ and $r_2 = |\vec{r}_2 - \vec{r}_3|$ are the relative distances. The space coordinates unit is $a_\mu = a_0/f$, where $a_0 = \hbar^2/(me^2) = 0.529\,177\,249(24)\text{\AA}$ is the Bohr radius. By this choice, the Hamiltonian is naturally expressed in reduced atomic units of energy $f\text{a.u.}$ ($= 13.605\,698\,1(40)\text{ eV} = 1\text{Ry}$ if $f = 0.5$), where $1\text{ a.u.} = e^2/a_0 = \alpha^2 mc^2$ is the atomic unit of energy and $\alpha = e^2/(\hbar c) = 0.007\,297\,353\,08(32)$ is the fine structure constant.

Approximate eigenfunctions of this Hamiltonian are obtained by using the variational method. The trial function is a finite linear combination

$$\begin{aligned} \Psi(\vec{r}_1, \vec{s}_1; \vec{r}_2, \vec{s}_2) = & \sum_{a,b,c,l_1,l_2} [q_{abc}^{l_1 l_2}(1, 2) \Phi_{abc\ l_1 l_2 LM}(\vec{r}_1, \vec{r}_2) \\ & + q_{abc}^{l_1 l_2}(2, 1) \Phi_{abc\ l_1 l_2 LM}(\vec{r}_2, \vec{r}_1)] \psi_{Sm_s}(\vec{s}_1, \vec{s}_2) \end{aligned} \quad (2)$$

of N_b basis elements $\Phi_{abc\ l_1 l_2 LM} \psi_{Sm_s}$. The orbital component $\Phi_{abc\ l_1 l_2 LM}$ is represented by the Hylleraas correlated wave function [1]

$$\Phi_{abc\ l_1 l_2 LM}(\vec{r}_1, \vec{r}_2) = r_1^a r_2^b r_{12}^c e^{-\alpha r_1 - \beta r_2} \mathcal{Y}_{LM}^{l_1 l_2}(1, 2) \quad (3)$$

which is a product between a polynomial in all relative radial variables and the orbital angular momentum eigenstates

$$\mathcal{Y}_{LM}^{l_1 l_2}(1, 2) = \sum_{m_1+m_2=M} C_{m_1 m_2 M}^{l_1 l_2 L} Y_{l_1 m_1}(\hat{r}_1) Y_{l_2 m_2}(\hat{r}_2) \quad , \quad (4)$$

where $\hat{r}_i = \vec{r}_i/r_i$, $i = 1, 2$, $\hat{r} = \vec{r}/r$ are unit vectors.

The spin function

$$\psi_{Sm_s} = \sum_{\mu_1+\mu_2=m_s} C_{\mu_1 \mu_2 m_s}^{\frac{1}{2} \frac{1}{2} S} |\frac{1}{2} \mu_1\rangle |\frac{1}{2} \mu_2\rangle \quad (5)$$

corresponds to singlet ($S = 0$) or triplet ($S = 1$) configurations, when the orbital part is symmetric ($q_{abc}^{l_1 l_2}(1, 2) = q_{abc}^{l_1 l_2}(2, 1)$) or antisymmetric ($q_{abc}^{l_1 l_2}(1, 2) = -q_{abc}^{l_1 l_2}(2, 1)$), respectively. The expansion coefficients $q_{abc}^{l_1 l_2}(1, 2)$, and the non-linear parameters α , β , have been determined previously [17] by using the variational equations

$$\delta_{q,\alpha,\beta} \frac{\langle \Psi | H_0 | \Psi \rangle}{\langle \Psi | \Psi \rangle} = 0 \quad . \quad (6)$$

The kinetic energy operator for the electron 1 is $-\nabla_1^2/2$, and its effective action in the Hylleraas model space is given by

$$\begin{aligned} \nabla_1^2 \Phi = & \left[\frac{1}{r_1^2} \frac{\partial}{\partial r_1} r_1^2 \frac{\partial}{\partial r_1} + \frac{1}{r^2} \frac{\partial}{\partial r} r^2 \frac{\partial}{\partial r} - \frac{\vec{l}_1^2}{r_1^2} \right. \\ & \left. + \frac{2(r_1 - r_2 \hat{r}_1 \cdot \hat{r}_2)}{r} \frac{\partial^2}{\partial r_1 \partial r} - \frac{2}{r_1 r} \vec{r}_2 \cdot \nabla_1^Y \frac{\partial}{\partial r} \right] \Phi \end{aligned} \quad (7)$$

where $\vec{l}_1 = \vec{r}_1 \times \vec{p}_1$ and $\nabla_1^Y = -i\hat{r}_1 \times \vec{l}_1$. A similar expression, obtained by permuting the indices 1 and 2, yields $\nabla_2^2 \Phi$.

The action of such operators on the Hylleraas basis states is complicated, and in the case of their product becomes tedious, involving hundreds of polynomial terms $r_1^a r_2^b r^c$, spherical harmonics, and singular delta functions. Therefore, in this work the matrix elements have been calculated by using a new procedure, based on the representation of the physical operators as linear combinations within a set $\mathcal{E} = \{E_k, k = 1, 2, \dots, n\}$ of n elementary operators. Although $\mathcal{E}$ is not a Lie algebra, this approach makes the numerical calculation more flexible, because the components E_k can be programmed individually, and they can be assembled as needed to form various complicated operators. By a suitable choice, the set $\mathcal{E}$ may account for several physical operators of interest. Some of the elementary operators used in the present calculation are presented in Appendix, Table I.

The accuracy of the wave function improves with the dimension N_b of the basis set. When N_b increases, the expectation value of the Hamiltonian $\langle H_0 \rangle_{(N_b)} = \langle \Psi | H_0 | \Psi \rangle$ decreases, and in principle, at the limit $N_b \rightarrow \infty$ the series $\langle H_0 \rangle_{(N_b)}$ approaches the exact ground state energy. Thus, a measure of the accuracy is provided by the deviation from 0, the known limit value, of the "convergence speed" $[\langle H_0 \rangle_{(N_b'')} - \langle H_0 \rangle_{(N_b')}] / (N_b'' - N_b')$, where N_b' and N_b'' are two consecutive values of N_b . A different measure of the accuracy is the variance $\sigma = \sqrt{\langle H_0^2 \rangle_{(N_b)} - \langle H_0 \rangle_{(N_b)}^2}$. The variational ground state energy $E_g = \langle H_0 \rangle_{(N_b)}$ [17] and the present results obtained for σ^2 are given in Appendix, Table II. The effective value defined in the Appendix-(b) is $E_g^{eff} = -0.524\,010\,140\,465\,960\,215\,38(56)f$ a.u.. Previous estimates of E_g in Ps^- by using the correlation-function hyperspherical-harmonic method and the stochastic variational method are $-0.524\,010\,139\,0 f$ a.u., respectively $-0.524\,010\,140\,452 f$ a.u. [16]. The effective value obtained in [15] by

using the extrapolation formula $E_g(N_b) = E_g^{eff} + A/N_b^p$ is $-0.524\ 010\ 140\ 465\ 956(8)\ f\ \text{a.u.}$.

III. Relativistic and QED Corrections

The quantum description of a relativistic charged fermion based on Dirac equation requires two spin-1/2 wave functions, $|\psi_+\rangle$ and $|\psi_-\rangle$, corresponding to the retarded and advanced waves, respectively. For the particle eigenstates with energy $E \sim mc^2$, $|\psi_-\rangle \sim K|\psi_+\rangle$, $K = \vec{\sigma} \cdot \vec{p}/(2mc)$, and the normalization condition $\langle\psi_+|\psi_+\rangle + \langle\psi_-|\psi_-\rangle = 1$, can be written in terms of the large components as $\langle\phi|\phi\rangle = 1$, where $|\phi\rangle = (1 + K^2/2)|\psi_+\rangle$. The antiparticle eigenstates are related in principle to the solutions with $E \sim -mc^2$, when $|\psi_-\rangle$ become the large components. Therefore, in general, the normalized eigenstate of a nonrelativistic Schrödinger equation for a particle or antiparticle should be seen as an approximation for $|\phi\rangle \approx (1 + K^2/2)|\psi_L\rangle$, where $|\psi_L\rangle$ denotes the corresponding large component. By definition, if $H_L|\psi_L\rangle = E_L|\psi_L\rangle$, then $H_{L,K}|\phi\rangle = E_L|\phi\rangle$, where $H_{L,K} = H_L + [K^2, H_L]/2$ includes both the nonrelativistic Hamiltonian and the lowest order correction terms.

In the interacting $e^-e^+e^-$ system $H_{L,K}$ will be restricted to $H_0 + H_1 + H_2$, where H_0 is given by Eq. (1). The term

$$H_1 = -\frac{1}{8m^3c^2}(p_1^4 + p_2^4 + p_3^4) \quad (8)$$

($p_i = |\vec{p}_i|$) takes into account the relativistic variation of the mass with velocity, and

$$H_2 = -\frac{1}{8m^2c^2} \sum_{i=1}^3 [\vec{p}_i \cdot, [\vec{p}_i, V]] \quad (9)$$

derives from the sum between the term $2(\vec{p}V) \cdot \vec{p}/(2mc)^2$ of H_L and the commutator $[K^2, V]/2 = 2[p^2, V]/(4mc)^2$. Here $V(R, R_1, R_2) = e^2(1/R - 1/R_1 - 1/R_2) \equiv \tilde{V}f\ \text{a.u.}$ and $R = a_\mu r$, $R_{1,2} = a_\mu r_{1,2}$.

The magnetic current-current interaction plus the retardation correction corresponding to the lowest-order Breit interaction are described by the additional term

$$M_2 = -\frac{e^2}{2m^2c^2} \{ R^{-1}[\vec{p}_1 \cdot \vec{p}_2 + \hat{r} \cdot (\hat{r} \cdot \vec{p}_1)\vec{p}_2] \\ - R_1^{-1}[\vec{p}_1 \cdot \vec{p}_3 + \hat{r}_1 \cdot (\hat{r}_1 \cdot \vec{p}_1)\vec{p}_3] - R_2^{-1}[\vec{p}_2 \cdot \vec{p}_3 + \hat{r}_2 \cdot (\hat{r}_2 \cdot \vec{p}_2)\vec{p}_3] \} \quad (10)$$

such that the total Hamiltonian for Ps^- that will be considered here is $H = H_0 + H_1 + H_2 + M_2$.

In a classical relativistic many-body system, the dynamical CM defined by the condition $\sum_i \vec{p}_i = 0$, is not necessarily the same as the geometrical centre of mass, located at $\vec{R}_{CM} = \sum_i m_i \vec{R}_i / \sum_i m_i$. In the present case, if $H = H_0 + H_1 + H_2 + M_2$, only the dynamical CM is inertial, because $[\vec{p}_1 + \vec{p}_2 + \vec{p}_3, H] = 0$, while $[\vec{r}_1 + \vec{r}_2 + \vec{r}_3, H]$ is not a constant. The use of the Hylleraas basis ensures that the present calculation takes place in the dynamical CM frame, because by the choice of the coordinates $-i\nabla_i = \vec{p}_i/(f\alpha mc)$ and $\nabla_3\Phi = (-\nabla_1 - \nabla_2)\Phi$ for any variational wavefunction Φ .

The expectation value $\langle H_1 \rangle = -(1/64)\langle \nabla_1^4 + \nabla_2^4 + \nabla_3^4 \rangle \alpha^2 f$ a.u. can be calculated either directly, or by assuming that in the ground state $\langle H_0 \mathcal{O}_p \rangle = \langle \mathcal{O}_p H_0 \rangle = E_g \langle \mathcal{O}_p \rangle$ for any operator $\mathcal{O}_p$, and using the equalities $\nabla_1^2 + \nabla_2^2 = 2(\tilde{H}_0 - \tilde{V} + f\nabla_1 \cdot \nabla_2)$, with $\tilde{H}_0 = H_0/(f\text{a.u.})$, and

$$\nabla_1^4 + \nabla_2^4 = 4(\tilde{H}_0 - \tilde{V} + f\nabla_1 \cdot \nabla_2)^2 - 2\nabla_1^2 \nabla_2^2, \quad (11)$$

$$\begin{aligned} \nabla_3^4 &= (-\vec{\nabla}_1 - \vec{\nabla}_2)^4 = \nabla_1^4 + \nabla_2^4 + 4(\nabla_1 \cdot \nabla_2)^2 \\ &\quad + 2\nabla_1^2 \nabla_2^2 + 4(\nabla_1^2 + \nabla_2^2)\nabla_1 \cdot \nabla_2. \end{aligned} \quad (12)$$

Although formally the same, within a finite basis the two expressions give slightly different results ($\langle \nabla_1^4 \rangle, \langle \nabla_1^4 \rangle_E$), presented in the Appendix, Table III. In the numerical estimates we have used only $\langle \nabla_1^4 \rangle_E$, because of its rapid convergence and higher accuracy in the effective value.

The term H_2 contains the singular operators $\Delta_1 \tilde{V} = -4\pi[\delta(\vec{r}) - \delta(\vec{r}_1)]$, $\Delta_2 \tilde{V} = -4\pi[\delta(\vec{r}) - \delta(\vec{r}_2)]$, and $\Delta_3 \tilde{V} = 4\pi[\delta(\vec{r}_1) + \delta(\vec{r}_2)]$, which yield

$$\langle H_2 \rangle = \alpha^2 \pi \langle \delta(\vec{r}_1) + \delta(\vec{r}_2) - \delta(\vec{r}) \rangle f^3 \text{ a.u.} \quad (13)$$

Previous estimates of $\langle \delta(\vec{R}_1) \rangle$ ($= a_\mu^{-3} \langle \delta(\vec{r}_1) \rangle$) in Ps^- by using the correlation - function hyperspherical - harmonic method and the stochastic variational method are 0.020 733 14(6) a_0^{-3} , respectively 0.020 731 048 976 a_0^{-3} [16]. The same methods give for $\langle \delta(\vec{R}) \rangle$ the values 0.000 170 997(2) a_0^{-3} and 0.000 171 112 600 741 a_0^{-3} , respectively [16]. The results of the present calculation, in the same units (a_0^{-3}), are listed in Appendix, Table IV as a function of the dimension N_b of the basis set.

The expectation values which appear in the calculus of $\langle M_2 \rangle$, obtained when $N_b = 324$ are

$$u_{ee} = \langle r^{-1} \nabla_1 \cdot \nabla_2 \rangle = -0.008\,267\,646\,67, \quad$$

$$v_{ee} = \langle r^{-1} \hat{r} \cdot (\hat{r} \cdot \nabla_1) \nabla_2 \rangle = 0.019\,610\,925\,35$$

and for $i = 1, 2$

$$u_{ep} = \langle r_i^{-1} (\nabla_i \cdot \nabla_3) \rangle = 1.535\,434\,049\,31$$

$$v_{ep} = \langle r_i^{-1} \hat{r}_i \cdot (\hat{r}_i \cdot \nabla_3) \nabla_i \rangle = -0.555\,009\,821\,912 \quad .$$

In terms of these variables, $\langle M_2 \rangle = 0.5\alpha^2 w f^3$ a.u. with $w = u_{ee} + v_{ee} - 2u_{ep} - 2v_{ep} = -1.949\,505\,176\,125$. For the 2528-dimensional basis set $w = -1.949\,505\,250\,368$. The average of the last three consecutive values, obtained for $N_b = 1990, 2276$ and 2528 gives the effective matrix element $w^{eff} = -1.949\,505\,250\,368(1)$.

The effective sum of the spin independent relativistic correction terms $\langle H_1 \rangle$, $\langle H_2 \rangle$ and $\langle M_2 \rangle$ is $-0.145\,476\,184\,397(8)\alpha^2 f$ a.u. which decrease the Ps^- ground state energy to

$$E_{g^*}^{eff} = \langle H \rangle^{eff} = E_g^{eff} - 0.145\,476\,184\,397(8)\alpha^2 f \text{ a.u.} \quad . \quad (14)$$

The same calculations yield for the corrected ground state energy $E_{g^*}^0$ in neutral positronium $E_{g^*}^0 = -(0.5 + 5\alpha^2/32)f$ a.u.. However, for this relativistic two-body system the finite mass corrections to the energy provided by the one-body Dirac equation can be obtained exactly up to the order $\alpha^2 \text{Ry}$ by using the formula [20]

$$E_{(n,j,Z)} = \frac{1}{\alpha^2} [\eta - 1 - \frac{f}{4}(\eta - 1)^2] \quad f \text{ a.u.} \quad (15)$$

where $\eta = 1/\sqrt{1 + (Z\alpha)^2/(n - \nu)^2}$, $\nu = j + 1/2 - \sqrt{(j + 1/2)^2 - (Z\alpha)^2}$. The expansion $E_{(1,1/2,1)} \approx -(0.5 + 5\alpha^2/32)f$ a.u. reproduces $E_{g^*}^0$, showing that the relativistic corrections given by the expectation value of H are reliable.

Within QED the constituents of the three body-system $e^-e^+e^-$ cease to be "elementary", because they are subject not only to the mutual two-body Coulomb-Breit interaction, but are also coupled to the vacuum fluctuations of the electromagnetic field $\vec{A}$ [21]. The interaction terms accounting for this coupling are represented by an infinite series of increasingly complicated Feynman diagrams with closed photon lines. However, the complexity is increased recursively, by taking into account at each order three basic processes, represented by the anomalous magnetic moment (vertex) corrections,

electron self-mass and vacuum polarization diagrams.

Although formally complicated, the main effect of the coupling to the field degrees of freedom is simply a change in the charge and mass parameters e and m of the theory. This contribution has already been taken into account, because it is included in the measured values of e and m used to define the atomic unit of energy. Though, the QED corrections in the interacting three body-system $e^-e^+e^-$ are not the same as for the free particles, and the differences still need to be considered.

The vacuum polarization properties have been studied first by Heisenberg [22] and Uehling [23], showing that a given charge density $\rho(\vec{R})$ induces a polarization charge $\delta\rho(\vec{R}) = -(\alpha/15\pi)\lambda_0^2\nabla^2\rho(\vec{R})$, where $\lambda_0 = \hbar/mc$ is the Compton wavelength of the electron. The induced charge leads to deviations from the standard Coulomb interaction. Thus, the vacuum behaves as an inhomogeneous dielectric, in which the mutual potential energy between two point-like charges Z_1 and Z_2 is [23]

$$V(R) = \frac{Z_1 Z_2 e^2}{R} \left[1 - \frac{\alpha}{\pi} R U(R) \right] , \quad (16)$$

by $U(R)$ denoting the Uehling potential. This potential is singular at $R = 0$, falls off exponentially for $R > 0$, and satisfies the integral condition $\int d^3R U(R) = -4\pi\lambda_0^2/15$. Therefore, it can be well approximated by a delta function, $U(R) = -4\pi(\lambda_0^2/15)\delta(\vec{R})$. In the case of Ps^- , the correction term introduced by this potential is

$$\langle H_{vp} \rangle = \frac{4}{15} \alpha^3 \langle \delta(\vec{r}) - \delta(\vec{r}_1) - \delta(\vec{r}_2) \rangle f^3 \text{ a.u.} \quad (17)$$

By using the effective values given in Appendix, Table IV, the contribution of the vacuum polarization to the Ps^- ground state energy is $\langle H_{vp} \rangle^{eff} = -0.022\,024\,212\,934\,6(7)\alpha^3 f \text{ a.u.}$. It is important to remark that this value takes into account the positron recoil (the "mass polarization" term) because the wave functions are obtained by minimizing the full nonrelativistic Hamiltonian. In neutral positronium, $\langle \delta(\vec{R}_1) \rangle_{Ps} = 1/(\pi a_\mu^3) = 1/(8\pi a_0^3)$, and the vacuum polarization correction is $-1/(15\pi) \alpha^3 f \text{ a.u.}$

As it was shown early by the Lamb shift measurements [24], the main QED correction appears however from the coupling to the vacuum fluctuations of the field rather than from the vacuum polarization ([21] p. 59). For

a free electron the ground state energy is given by its rest mass $m = m_b + \delta m$, consisting of the uncoupled value m_b and the positive renormalization term $\delta m = (3\alpha m_b/2\pi) \ln(\Lambda/m_b)$ due to the electromagnetic self-energy, where Λ is a large (formally infinite) cutoff mass.

Similarly, the coupling to the field modes also affects the intrinsic excitations of a many-body system. In a bounded N -particle system, the shift ΔE_n in the energy $E_n = \langle n|H_0|n\rangle$ of the level $|n\rangle$ due to the exchange of a transverse photon can be obtained by using the time-independent second-order perturbation expression

$$\Delta E_n = -\langle n, 0_f | H_c(\vec{A}) \frac{1}{H_0 + H_A - E_n} H_c(\vec{A}) | n, 0_f \rangle \quad (18)$$

Here $H_c(\vec{A}) = \sum_{i=1}^N h_i(\vec{A})$ is the sum over all particles of the one-body coupling terms $h_i(\vec{A}) = -e_i \vec{\alpha}_i \cdot \vec{A}_{(\vec{R}_i)}$, $\vec{\alpha}_i \approx \vec{p}_i/(m_i c)$,

$$\vec{A}_{(\vec{r})} = \frac{\sqrt{\hbar c}}{2\pi} \int \frac{d^3 k}{\sqrt{k}} \sum_{\lambda=1,2} \vec{\epsilon}_\lambda (\hat{a}_{k\lambda}^\dagger e^{-i\vec{k}\cdot\vec{r}} + \hat{a}_{k\lambda} e^{i\vec{k}\cdot\vec{r}}) \quad (19)$$

is the quantized transverse vector potential of the photon ($\vec{\epsilon}_\lambda \cdot \vec{k} = 0$, $\epsilon_\lambda^2 = 1$), $H_A = \int d^3 k \sum_{\lambda=1,2} \hbar c k a_{k\lambda}^\dagger a_{k\lambda}$ is the free field Hamiltonian, and $|0_f\rangle$ denotes the photon vacuum. This shift has the form $\Delta E_n = \sum_{i=1}^N X_i^n + \sum_{i<j} Y_{ij}^n$, where

$$X_i^n = -\langle n, 0_f | h_i(\vec{A}) \frac{1}{H_0 + H_A - E_n} h_i(\vec{A}) | n, 0_f \rangle \quad (20)$$

and

$$Y_{ij}^n = -2\text{Re}[\langle n, 0_f | h_i(\vec{A}) \frac{1}{H_0 + H_A - E_n} h_j(\vec{A}) | n, 0_f \rangle] \quad (21)$$

It is important to remark that the interaction with the vacuum field fluctuations may affect not only the intrinsic dynamics, but also the center of mass. In a classical two-body system coupled to the field, H_c can be written in terms of the canonical pairs $(\vec{r}, \vec{p}_\mu) \equiv (\vec{R}_1 - \vec{R}_2, \mu\vec{p}_1/m_1 - \mu\vec{p}_2/m_2)$ and $(\vec{R}_{CM}, \vec{P}_{CM}) \equiv (\mu\vec{R}_1/m_2 - \mu\vec{R}_2/m_1, \vec{p}_1 + \vec{p}_2)$ of intrinsic and respectively, center of mass variables, as

$$\vec{p}_\mu \cdot \left[\frac{e_2}{m_2} \vec{A}_{(R_2)} - \frac{e_1}{m_1} \vec{A}_{(R_1)} \right] - \frac{\mu}{m_1 m_2} \vec{P}_{CM} \cdot [e_1 \vec{A}_{(R_1)} + e_2 \vec{A}_{(R_2)}] \quad .$$

This expression shows that in a neutral two-body system (such as Ps) the center of mass energy is not affected by the field only if $\vec{A}_{(R_1)} = \vec{A}_{(R_2)}$, or when the size of the system is negligible compared to the photon wavelength (dipole approximation).

In a quantum N -body system it is convenient to take advantage of the finite size effects by writing $\vec{A}$ as the incoherent sum of long and short wavelength components, $\vec{A}_L$ and $\vec{A}_S$, obtained by decomposing $\int d^3k$ as $\int_{|\vec{k}| \leq k_L} d^3k + \int_{k_L < |\vec{k}| < k_M}$, where k_L and k_M are cutoff parameters. Each domain brings its own contribution to the matrix elements, which can be similarly decomposed as

$$X_i^n = X_i^{Ln} + X_i^{Sn} \quad , \quad Y_{ij}^n = Y_{ij}^{Ln} + Y_{ij}^{Sn} \quad . \quad (22)$$

At the end of the calculation k_L should disappear, while $k_M \rightarrow \infty$.

If H_0 consists of the kinetic energy term plus a local potential V , then a non-relativistic calculation within the dipole approximation yields

$$X_i^{Ln} = -\frac{\alpha}{3\pi m^2 c^2} [2\hbar c k_L \langle n | \vec{p}_i^2 | n \rangle + \langle n | [\vec{p}_i, [\vec{p}_i, V]] | n \rangle \ln \frac{k_L}{k_R} - 2B_{ii}^n] \quad , \quad (23)$$

where $k_R = R_M/\hbar c$, R_M is a dimensional constant with units of energy, and B_{ii}^n are the diagonal elements of the matrix $[B_{ij}^n]$ defined by

$$B_{ij}^n = \sum_m (E_n - E_m) \text{Re}(\langle n | \vec{p}_i | m \rangle \cdot \langle m | \vec{p}_j | n \rangle) \ln \frac{|E_n - E_m|}{R_M} \quad . \quad (24)$$

The first term depends only on the kinetic energy, and it can be written as $-\delta m_L \langle n | \vec{p}_i^2 | n \rangle / (2m^2)$, $\delta m_L = 4mr_e k_L / (3\pi)$, where $r_e = \alpha\hbar/(mc)$ denotes the classical radius of the electron. It contributes also to the energy of a free particle ($V = 0$) and has the structure of a first-order perturbation shift induced by a variation δm_L of the nonrelativistic mass. Thus, such terms can be taken into account simply by a redefinition of the cutoff mass Λ .

A relativistic calculation of the one-body QED correction arising from the exchange of a transverse hard photon at a Coulomb vertex [25], [21] p.177, yields

$$X_i^{Sn} = \frac{\alpha\hbar^2}{3\pi m^2 c^2} \left(\ln \frac{mc}{2\hbar k_L} + \frac{5}{6} \right) \langle n | \Delta_i V | n \rangle \quad , \quad (25)$$

($5/6 = 11/24 + 3/8$) such that

$$X_i^n = -\frac{\delta m_L}{2m^2} \langle n | \vec{p}_i^2 | n \rangle + \frac{\alpha}{3\pi m^2 c^2} \left[\hbar^2 \left(\ln \frac{mc}{2\hbar k_R} + \frac{5}{6} \right) \langle \Delta_i V \rangle_n + 2B_{ii}^n \right] \quad . \quad (26)$$

The quantity Y_{ij}^{Ln} can be expressed as

$$Y_{ij}^{Ln} = -\frac{\delta m_L}{m^2} \frac{e_i e_j}{e^2} \langle \vec{p}_i \cdot \vec{p}_j \rangle_n - \frac{2\alpha}{3\pi m^2 c^2} \frac{e_i e_j}{e^2} \{ \langle [\vec{p}_i \cdot, [\vec{p}_j, V]] \rangle_n \ln \frac{k_L}{k_R} - 2B_{ij}^n \} . \quad (27)$$

In the case of Ps^- there are three terms X_i^n , one for each electron ($i = 1, 2$) and one for the positron ($i = 3$), and three terms Y_{ij}^{Ln} , $i < j$. The contribution to ΔE_n arising from the terms linear in δm_L of X and Y^L is $\delta_0 E_n = -\delta m_L \langle n | (\vec{p}_1 + \vec{p}_2 - \vec{p}_3)^2 | n \rangle / (2m^2)$. In the dynamical CM frame, this energy shift can be accounted, for example, by an effective variation $3\delta m_L$ in the total mass of the electron-electron pair and δm_L in the mass of the positron, or by a variation of $4\delta m_L$ in only one of them.

The definition of the Bethe logarithm $\beta_n \equiv 2B_{33}^n / \langle [\vec{p}_3, \cdot [\vec{p}_3, H_0]] \rangle_n$, and the identity $m^2 \sum_{i,j} e_i e_j B^n(i, j) / (m_i m_j) = e^2 B^n(3, 3) (1 + m/m_3)^2$ (valid if $m_1 = m_2 = m$ and $\langle n | \sum_i \vec{p}_i | n' \rangle = 0$ for any n, n'), show that $\sum_i X_i^n + \sum_{i < j} Y_{ij}^{Ln} = \delta_0 E_n + \delta_1 E_n + \delta_{2L} E_n$, where

$$\begin{aligned} \delta_1 E_n &= \frac{\alpha \hbar^2}{3\pi m^2 c^2} [-4\beta_n \langle \Delta_3 V \rangle_n + (\ln \frac{mc}{2\hbar k_R} + \frac{5}{6}) \sum_{i=1}^3 \langle \Delta_i V \rangle_n] \quad (28) \\ &= \frac{4\alpha^3}{3} [-4\beta_n \langle \delta^3(r_1) + \delta^3(r_2) \rangle_n + 2(\ln \frac{mc}{2\hbar k_R} + \frac{5}{6}) \langle \delta^3(r_1) + \delta^3(r_2) - \delta^3(r) \rangle_n] f^3 \text{ a.u.} \\ \text{and } \delta_{2L} E_n &\equiv \sum_{i < j} \langle W_{ij}^L \rangle_n \text{ is given by the expectation value of the potential} \end{aligned}$$

$$W_{ij}^L(k_L) = \frac{8\alpha^3}{3} \ln \frac{k_L}{k_R} \delta^3(r_{ij}) f^3 \text{ a.u..} \quad (29)$$

The term Y_{ij}^{Sn} due to the exchange of a short wavelength (hard) transverse photon between different particles will be decomposed as $Y_{ij}^{Sn} = Y_{ij}^{S2n} + Y_{ij}^{S3n}$, according to the expansion $1/(H_0 + H_A - E_n) \approx 1/H_A - (H_0 - E_n)/(H_A)^2$. The contribution from $1/H_A$ is

$$Y_{ij}^{S2n} = -2 \langle n, 0_f | h_i(\vec{A}_S) H_A^{-1} h_j(\vec{A}_S) | n, 0_f \rangle . \quad (30)$$

In the limit $k_L \rightarrow 0$, $k_M \rightarrow \infty$, the integral over k in this matrix element can be evaluated by using the identity

$$\int \frac{d^3 k}{k^2} e^{i\vec{k} \cdot \vec{r}} (\vec{A} \cdot \vec{B} - \hat{k} \cdot \vec{A} \hat{k} \cdot \vec{B}) = \frac{\pi}{r} (\vec{A} \cdot \vec{B} + \hat{r} \cdot \vec{A} \hat{r} \cdot \vec{B}) ,$$

showing that the sum $\sum_{i<j} Y_{ij}^{S2n}$ becomes the two-body term $\langle M_2 \rangle_n$ of order $\alpha^2 \text{Ry}$, already taken into account. Thus, the only new contribution is the next-order term

$$Y_{ij}^{S3n} = 2\text{Re}[\langle n, 0_f | h_i(\vec{A}_S) \frac{H_0 - E_n}{H_A^2} h_j(\vec{A}_S) | n, 0_f \rangle] \quad (31)$$

which is the expectation value of the two-body potential

$$W_{ij}^S(k_L, k_M) = \frac{2\alpha^3}{3\pi} \left[\frac{3}{2} f(k_L, k_M, r_{ij}) + 4\pi \delta^3(r_{ij}) \ln \frac{k_M}{k_L} \right] f^3 \text{ a.u.} \quad (32)$$

Here $f(k_L, k_M, r) = 2[j_0(k_M r) + j_2(k_M r) - j_0(k_L r) - j_2(k_L r)]/(3r^3)$ is the function introduced by Araki [26], written in terms of the spherical Bessel functions j_0, j_2 . When $k_L \rightarrow 0$, $k_M \rightarrow \infty$, $f(0, \infty, r) = -2/(3r^3)$, but the logarithmic factor in the second term of W_{ij}^S is divergent at both limits. However, the divergence in k_L is cancelled by the low-energy term, and the sum $W_{ij}(k_M) = W_{ij}^L(k_L) + W_{ij}^S(k_L, k_M)$,

$$W_{ij}(k_M) = \frac{2\alpha^3}{3\pi} \left[\frac{3}{2} f(0, k_M, r_{ij}) + \frac{\delta(r_{ij})}{r_{ij}^2} \ln \frac{k_M}{k_R} \right] f^3 \text{ a.u.} , \quad (33)$$

is independent of k_L . The divergent factor containing k_M contributes only when $|n\rangle$ is an S state, but in this case the expectation value $\langle 1/r^3 \rangle_n$ is also logarithmically divergent. It is however possible to define a limit for the sum of these infinite terms in the sense of the principal value. Let

$$D(a, r) = \frac{\theta(r-a)}{r^3} - \frac{\delta(r)}{r^2} \ln \frac{a_\mu}{a} \quad (34)$$

be a distribution depending on the positive radius parameter $a = \eta/k_M$, where η is a positive scale factor. Because $r^2 \partial_a D(a, r) = [\delta(r) - \delta(r-a)]/a$, when $k_M \rightarrow \infty$ the expectation value $\langle D(a, r) \rangle_n$ is finite. In terms of this distribution we can define the principal value

$$\mathcal{P} \left[\frac{3}{2} f(0, k_M, r) + \frac{\delta(r)}{r^2} \ln \frac{k_M}{k_R} \right] |_{k_M \rightarrow \infty} = 4\pi \delta^3(r) \ln \frac{\eta}{a_\mu k_R} - \lim_{a \rightarrow 0} D(a, r) . \quad (35)$$

The choice of a scale factor $\eta = e^{\frac{4}{3}-\gamma}$, where γ is the Euler's constant, yields the formula used by Araki [26]

$$\langle W_{ij} \rangle_n = -\frac{2\alpha^3}{3\pi} \{ Q_{ij}^n + 4\pi \langle \delta^3(r_{ij}) \rangle_n [\ln a_\mu k_R - \frac{4}{3}] \} f^3 \text{ a.u.} , \quad (36)$$

where $Q_{ij}^n = \lim_{a \rightarrow 0} \langle D(a, r) + 4\pi\gamma\delta^3(r_{ij}) \rangle_n$. In Ps^- this yields for the effective two-body contribution $\delta_2 E_n = \delta_{2L} E_n + \sum_{i < j} \langle W_{ij}^S \rangle_n = \sum_{i < j} \langle W_{ij} \rangle_n$ the expression

$$\begin{aligned} \delta_2 E_n = & -\frac{2\alpha^3}{3\pi} [Q_{12}^n + Q_{13}^n + Q_{23}^n] \\ & + 4\pi(\ln a_\mu k_R - \frac{4}{3}) \langle n | \delta^3(r_1) + \delta^3(r_2) + \delta^3(r) | n \rangle] f^3 \text{ a.u.} . \end{aligned} \quad (37)$$

Summarizing these results, the effective QED contribution of order α^3 to the energy level E_n of Ps^- due to the exchange of a transverse photon is $\delta_{1p} E_n = \delta_1 E_n + \delta_2 E_n$. This sum is independent of the arbitrary energy unit R_M , as it should, but for the purpose of numerical calculations we choose $R_M = f\text{Ry}$. With this choice, $a_\mu k_R = \alpha/2$, and $mc/(2\hbar k_R) = 1/(f\alpha^2)$.

The corresponding terms for positronium can be obtained from the expressions given above simply by neglecting all the expectation values containing the variables r_2 and r , involving the second electron. For the Ps ground state $Q_{13}^{Ps} = -4 \ln 2$, while β_g^{Ps} is the same as the Bethe logarithm for hydrogen $\beta_g^H = 2.984\,128\,555\,765\,497\,611(4)$, each Bethe logarithm being calculated using the corresponding reduced Rydberg constant [27]. For the Ps^- ground state the numerical values of Q_{12}^g and Q_{13}^g used in the present estimates are listed in Table V, while $\beta_g = 3.005\,030(2)$ [27] (including the finite mass correction).

To the same order we should consider also the double photon exchange term (including the Coulomb part) $\delta_{2p} E_n$ [26] [28]

$$\begin{aligned} \delta_{2p} E_n = & -\frac{\alpha^3}{2\pi} [Q_{12}^n + Q_{13}^n + Q_{23}^n] \\ & - 4\pi(\ln f\alpha - \frac{4}{3} \ln 2 + \frac{13}{6}) \langle n | \delta^3(r_1) + \delta^3(r_2) + \delta^3(r) | n \rangle] f^3 \text{ a.u.} , \end{aligned} \quad (38)$$

and the energy shift associated with the two-photon decay.

In general, any coupling which makes the levels unstable produces a complex energy shift $\Delta_c E_n = \delta_c E_n - i\Gamma_c^n/2$, where $\delta_c E_n$ is a correction to the level centroid, $\lambda_c^n = \Gamma_c^n/\hbar$ is the decay rate, and c denotes the decay channel. Neutral positronium normally decays by spontaneous e^+e^- annihilation into two photons if the total spin $S_{ep} = 0$, ($\vec{S}_{ep} = \vec{s}_e + \vec{s}_p$), and in three photons if $S_{ep} = 1$ [29]. The corresponding decay rates are such that $\Gamma_{3\gamma} \sim \alpha\Gamma_{2\gamma}$,

and the first correction arises from two photon annihilation. In this channel $\delta_{2\gamma}E_n/\Gamma_{2\gamma}^n = -(1 - \ln 2)/\pi$ [30], where

$$\Gamma_{2\gamma}^n = 2\pi\alpha^3 \langle (2 - \vec{S}_{ep}^2) \delta^3(r_1) \rangle_n f^3 \text{ a.u.} . \quad (39)$$

For the Ps ground state ($\vec{S}_{ep} = 0$) this yields a decay rate $\lambda_{(Ps, 2\gamma)} = \alpha^3 \text{Ry}/\hbar = 8.04 \text{nsec}^{-1}$, in good agreement with the experimental result $7.99(11) \text{nsec}^{-1}$ [31].

In the Ps^- ground state the electron spins are coupled to 0, and the two-photon annihilation can take place between the positron and any of the two electrons. The total rate depends on $\langle \vec{S}_{13}^2 + \vec{S}_{23}^2 \rangle_g = 3$, and can be expressed in the form

$$\Gamma_{(Ps^-, 2\gamma)} = 2\pi\alpha^3 \langle \delta^3(r_1) \rangle f^3 \text{ a.u.} . \quad (40)$$

The effective ground state expectation value $\langle \delta(\vec{R}_1/a_0) \rangle$ given in the Appendix, Table IV, yields $\lambda_{(Ps^-, 2\gamma)} = 2.092\,797(1) \text{ nsec}^{-1}$, in good agreement with the previous estimates [32] and the experimental result $2.09(9) \text{ nsec}^{-1}$ [12]. The corresponding level shift is $\delta_{2\gamma}E_g = -(1 - \ln 2)\Gamma_{(Ps^-, 2\gamma)}/\pi$.

Summarizing the results of these calculations, the effective ground state expectation values of the nonrelativistic Hamiltonian and the first relativistic and QED corrections for Ps and Ps^- are collected in the following table

$E[\text{Ry}]$	Ps	Ps^-	$E_{\text{Ps}} - E_{\text{Ps}^-}$
E_g	$-1/2$	$-0.524\,010\,140\,465\,960\,215\,38$ $\pm 0.56 \times 10^{-18}$	$0.024\,010\,140\,465\,960\,215\,38$ $\pm 0.56 \times 10^{-18}$
$\langle H_1 \rangle / \alpha^2$	$-5/32$	$-0.161\,254\,673\,938\,50(6)$	$0.005\,004\,673\,938\,50(6)$
$\langle H_2 \rangle / \alpha^2$	$1/4$	$0.259\,466\,645\,837(8)$	$-0.009\,466\,645\,837(8)$
$\langle M_2 \rangle / \alpha^2$	$-1/4$	$-0.243\,688\,156\,296\,0(1)$	$-0.006\,311\,843\,704\,0(1)$
$\langle H_{vp} \rangle / \alpha^3$	$-1/(15\pi)$	$-0.022\,024\,212\,934\,6(7)$	$0.000\,803\,553\,855\,7(7)$
$\delta_{1p}E_g/\alpha^3$	$2.766\,873\,00(3)$	$3.006\,491\,9(9)$	$-0.239\,618\,9(9)$
$\delta_{2p}E_g/\alpha^3$	$-0.585\,335\,778(7)$	$-0.510\,831\,605(7)$	$-0.074\,504\,17(1)$
$\delta_{2\gamma}E_g/\alpha^3$	$-(1 - \ln 2)/\pi$	$-0.025\,448\,161\,055(1)$	$-0.072\,226\,124\,976(1)$

In Ps^- the nonrelativistic one electron binding energy $0.024\,010\,140\,465\,960\,215\,38(56) \text{ Ry}$ is practically the same as the one determined in [15], and close to the older estimate of $0.024\,010\,113 \text{ Ry}$ [32]. The effect of the correction terms is to decrease slightly this energy to

$$B' = [0.024\,010\,140\,465\,960\,215\,38(56) - 0.010\,773\,815\,602(8)\alpha^2] \quad (41)$$

$$-0.385\,545\,7(9)\alpha^3] \text{ Ry} = 0.024\,009\,416\,924\,85(6) \text{ Ry} .$$

In positronium the observed hyperfine splitting of $1.160\,963(9)\alpha^2 \text{ Ry}$ between the otherwise degenerate ground state components corresponds to an additional spin-spin contact term [30]

$$\delta_s^{ep} E_n^0 = 2\pi\alpha^2 \langle \delta^3(r_1) \{ \frac{4}{3} \vec{s}_e \cdot \vec{s}_p (1 - \frac{\alpha}{2\pi}) + \frac{1}{2} \vec{S}_{ep}^2 [1 - (\frac{26}{9} + \ln 4) \frac{\alpha}{\pi}] \} \rangle_n f^3 \text{ a.u.} , \quad (42)$$

shifting the energy of the singlet by $\delta_s^{ep} E_g^0 = -2\alpha^2(1 - \alpha/2\pi)f^3 \text{ a.u.} = -0.265\,947\,576(23) \times 10^{-4} \text{ Ry}$. In Ps^- the electron-electron spin-spin dependent energy shift is [26]

$$\delta_s^{ee} E_g = -\frac{8\pi}{3} \alpha^2 (1 + \frac{5}{2\pi} \alpha) \langle \vec{s}_1 \cdot \vec{s}_2 \delta^3(r) \rangle f^3 \text{ a.u.} , \quad (43)$$

while the two electron-positron spin-spin contact terms contribute by

$$\delta_s^{2ep} E_g = \pi\alpha^2 \langle \delta^3(r_1) (\vec{S}_{13}^2 + \vec{S}_{23}^2) [1 - (\frac{26}{9} + \ln 4) \frac{\alpha}{\pi}] \rangle f^3 \text{ a.u.} . \quad (44)$$

These formulas yield an additional shift of the Ps^- ground state energy $\delta_s E_g = \delta_s^{ee} E_g + \delta_s^{2ep} E_g = 0.207\,196\,744(18) \times 10^{-4} \text{ Ry}$, and a contribution to its binding energy of $\delta_s^{ep} E_g^0 - \delta_s E_g = -0.473\,144\,32(3) \times 10^{-4} \text{ Ry}$. Including the spin-spin contact terms, the binding energy becomes

$$B'' = B' + \delta_s^{ep} E_g^0 - \delta_s E_g = 0.023\,962\,102\,492(3) \text{ Ry} . \quad (45)$$

IV. Summary and Conclusions

The calculation of the relativistic and QED corrections to the energy levels of a quantum three-body system represents a challenging problem the modern theory. Difficulties appear both at conceptual and computational levels, as there is no satisfactory relativistic many-body quantum theory, and the nonrelativistic problem is not integrable.

A quantum three-body system thoroughly investigated since the early days of quantum mechanics is the helium atom. In this system a major simplification occurs, because the reduced electron mass μ is smaller than the

mass of the positive charge by a factor 1.3707×10^{-4} , and to a first approximation the relative motion of the nucleus in the center of mass frame can be neglected. The relativistic invariance is partly restored by the Breit interaction, and highly accurate nonrelativistic wave functions can be obtained numerically, from variational calculations. Within this framework, a perturbative treatment of the relativistic and QED corrections gives energy levels in remarkable agreement with experiment [3] [4].

The same procedure was applied in this work to the negative positronium ion. However, by contrast to helium, all three particles have equal mass, and a perturbative treatment of the positron motion becomes inappropriate.

The accuracy of the nonrelativistic energy and variational ground state wave function was discussed in Sect. II. The effective value of E_g estimated here is $-0.262\,005\,070\,232\,980\,107\,69(28)$ a.u., the same as in [15], [17] and close within 10^{-8} to estimates obtained by other methods [16]. The variance of the Hamiltonian for the largest (2528-dimensional) basis set is 2.78×10^{-8} a.u., smaller than the level width $\Gamma = \hbar\lambda = 5.06 \times 10^{-8}$ a.u. due to the $(2\gamma)e^-$ decay.

The calculation of the first relativistic and QED corrections has been presented in Sect. III. Formal expressions of these correction terms have been known for a long time, but the mass polarization term, which cannot be neglected for Ps^- , increases dramatically the complexity of the numerical calculations. In this work the calculations have been performed by representing the action of the physical operators on the Hylleraas basis states in terms of a suitable set of elementary operators (Appendix, Table I). The numerical values obtained for some of the most important matrix elements are summarized in the Tables II-V of the Appendix. It was found that the spin independent relativistic terms contribute to the Ps^- ground state energy by $-0.072\,738\,092\,198(4)\alpha^2$ a.u. and the lowest order QED corrections by $1.224\,094\,00(44)\alpha^3$ a.u.. These terms decrease the ground state energy to $E_{g*}^{eff} = -0.262\,008\,467\,959\,9(4)$ a.u.. Both contributions decrease slightly also the one electron binding energy, from the nonrelativistic value $0.012\,005\,070\,232\,980\,10(3)$ a.u. to $0.012\,004\,708\,462\,43(3)$ a.u.. A much larger contribution appears however from the spin-dependent contact terms, which raise the ground state energy to $E_{g*}^{eff} = -0.261\,998\,108\,122(1)$ a.u., and further decrease the binding energy to $0.011\,981\,051\,246(2)$ a.u.. The calculated decay rate by two photon emission is $2.092\,797(1)$ nsec $^{-1}$, close to the previous theoretical results and to the measured value [12].

The binding and ground state energies are shifted also by relativistic corrections of order $\alpha^3\text{Ry}$ containing spin-orbit magnetic interactions. These contributions are also important, and should be included in the further attempts to improve the accuracy of the present estimates.

Appendix

(a) Elementary operators

In terms of the elementary operators defined in Table I, the nonrelativistic Hamiltonian contains linear combinations such as

$$\begin{aligned} \left(-\frac{1}{2}\nabla_1^2 - \frac{1}{2}\nabla_2^2\right)\Phi &= (E_2 + E_3 - E_4 + E_5 + E_6 - 0.5E_9 \\ &\quad - 0.5E_{10} - 0.5E_{11} - 0.5E_{12})\Phi \end{aligned}$$

for the kinetic energy part, $\tilde{V}\Phi = -E_{13}\Phi$ for the potential energy and

$$\nabla_1 \cdot \nabla_2 \Phi = (E_1 + E_2 + E_3 - E_4 + E_5 + E_6 + E_7 + E_8)\Phi$$

for the recoil term. The action on the basis states of a linear combination $\hat{O}_p = \sum_k f_{pk} E_k$, has the general form

$$\hat{O}_p \Phi_{x_0} = \sum_{t=1}^{N_p} q_{px}^t A_t \Phi_{x_t}$$

where x denotes the whole set of indices of Φ , q_{px}^t is a factor determined by x and the action of E_k on the radial functions, while A_t denotes the remaining angular operator. In this sum the same state $A_t \Phi_{x_t}$ (or $A_t A_{t'} \Phi_{x_{tt'}}$ in the case of a product) may appear several times with different scalar factors q . Therefore, in general it is possible to reduce the number of terms from N_p to $N_r < N_p$ by partial summations, before the effective calculation of the matrix elements. The reduction increases the speed of the numerical calculation, because N_r can be significantly smaller than N_p . For example, $N_p : N_r$ is 20 : 18 in the case of $\vec{\nabla}_1 \cdot \vec{\nabla}_2$, 400 : 235 for $(\vec{\nabla}_1 \cdot \vec{\nabla}_2)^2$, while for H_0 and H_0^2 is 31 : 24 and 961 : 427, respectively.

In the computer program the action of E_k consists of both arithmetic and symbolic operations. The arithmetic operations are determined by its action on the radial function $r_1^a r_2^b r_{12}^c e^{-\alpha r_1 - \beta r_2}$. The number n_k of distinct polynomial terms generated by this action is given in the second column of Table I. The symbolic operation corresponds to the action of E_k on the spherical harmonics in $\mathcal{Y}_{LM}^{l_1 l_2}$, and is coded by a character denoting angular operators such as $\hat{r}_1 \cdot \hat{r}_2$ or $\nabla_1^Y \cdot \nabla_2^Y$.

(b) *Error estimates*

The series of numerical values presented in Tables II-V appear to be convergent, but for comparison with experiment, it is useful to provide also a single effective value, representing the expected result of the present calculation when $N_b \rightarrow \infty$. The procedure adopted here to define this value depends on the manner of convergence. In the case of a sequence $\{f_n\}$ convergent as an alternating series, the effective value $f_{eff} \pm \sigma_f$, given in the last row, was defined as the arithmetic average of its last three consecutive terms, by $f_{eff} = (f_{n_x} + f_{n_y} + f_{n_z})/3$, $n_x < n_y < n_z$, and $\sigma_f^2 = [(f_{n_x} - f_{eff})^2 + (f_{n_y} - f_{eff})^2 + (f_{n_z} - f_{eff})^2]/3$. If $\{f_n\}$ approaches the limit by monotonous increase or decrease, then we can assume that the series can be extended to infinity by the function $F(n) = f_{eff} + Ae^{-\gamma n}$. The matching equations $F(n_x) = f_{n_x}$, $F(n_y) = f_{n_y}$, $F(n_z) = f_{n_z}$ between $F(n)$ and the last three calculated numerical values yield the parameter f_{eff} in the form [17]

$$f_{eff} = f_{n_y} + \frac{f_{n_y} - f_{n_x}}{R - 1} .$$

Here $R \equiv e^{\gamma(n_y - n_x)}$ is the solution of the equation $R - 1 = R_y[1 - R^{(n_y - n_z)/(n_y - n_x)}]$, where

$$R_y = \frac{f_{n_y} - f_{n_x}}{f_{n_z} - f_{n_y}} .$$

The error is assumed to be

$$\sigma_f = |f_{n_y} - f_{eff}| = \left| \frac{f_{n_y} - f_{n_x}}{R - 1} \right| .$$

If $n_y - n_x = n_z - n_y$, then $R = R_y$. When n is simply N_b , then $n_y - n_x = 286$ is larger, but close to $n_z - n_y = 252$, and $R = R_y$ still provides a reasonable

estimate.

(c) *Expectation values of p^4*

The expectation values $\langle \nabla_1^4 \rangle = \langle \nabla_2^4 \rangle$ for electrons in the Ps^- ground state can be calculated numerically either directly, as

$$\langle \nabla_1^4 \rangle = 4\langle (E_2 - 0.5E_4 + E_6 - 0.5E_9 - 0.5E_{11})^2 \rangle$$

or as $\langle \nabla_1^4 \rangle_E = 0.5\langle \nabla_1^4 + \nabla_2^4 \rangle_E = 0.5\langle (\nabla_1^2 + \nabla_2^2)^2 \rangle_E - \langle \nabla_1^2 \nabla_2^2 \rangle$, where the first term is expressed in the form

$$\begin{aligned} 0.5\langle (\nabla_1^2 + \nabla_2^2)^2 \rangle_E &= 2\langle (\tilde{E}_g - \tilde{V} + f\nabla_1 \cdot \nabla_2)^2 \rangle \\ &= 2[\tilde{E}_g^2 - 2\tilde{E}_g\langle \tilde{V} - f\nabla_1 \cdot \nabla_2 \rangle + \langle (\tilde{V} - f\nabla_1 \cdot \nabla_2)^2 \rangle] \end{aligned}$$

($\tilde{E}_g \equiv E_g/f$ a.u.) by assuming that the variational ground state is practically eigenstate of H_0 . Although formally the same at the limit $N_b \rightarrow \infty$, the numerical values obtained for $\langle \nabla_1^4 \rangle$ and $\langle \nabla_1^4 \rangle_E$ at finite N_b are slightly different. These estimates are given as a function of the basis size N_b in the first two columns of Table III. The third column contains the relativistic correction term for the positron, given by

$$\langle \nabla_3^4 \rangle_E = \langle (\nabla_1 + \nabla_2)^4 \rangle_E = 2\langle \nabla_1^4 \rangle_E + 2\langle \nabla_1^2 \nabla_2^2 \rangle - 8\langle (\tilde{E}_g - \tilde{V})\nabla_1 \cdot \nabla_2 \rangle \quad .$$

(d) *QED corrections of order α^3 in the limit $m_3 \rightarrow \infty$*

When $m_3 \rightarrow \infty$ the vacuum polarization term and the contribution of the electron-electron spin dependent contact interaction $\delta_s^{ee} E_n|_{\alpha^3} = -(20/3)\langle \vec{s}_1 \cdot \vec{s}_2 \delta^3(r) \rangle_n \alpha^3 f^3$ a.u. remain the same, but $X_3^n = 0$, $Y_{i3}^n = 0$, and $\delta_{1p} E_n$ becomes

$$\begin{aligned} \delta_{1p} E_n^\infty &= \alpha^3 \left[\frac{4}{3} \left(\ln \frac{mc}{2\hbar k_R} + \frac{5}{6} - \beta_n \right) \langle \delta^3(r_1) + \delta^3(r_2) \rangle_n \right. \\ &\quad \left. - \frac{8}{3} \left(\ln \frac{mc}{2\hbar k_R} - \frac{1}{2} + \ln a_\mu k_R \right) \langle \delta^3(r) \rangle_n - \frac{2}{3\pi} Q_{12}^n \right] f^3 \text{ a.u.} \quad . \end{aligned}$$

The two-photon contribution reduces to

$$\delta_{2p} E_n^\infty = \alpha^3 \left[-\frac{Q_{12}^n}{2\pi} + 2 \left(\ln f\alpha - \frac{4}{3} \ln 2 + \frac{13}{6} \right) \langle \delta^3(r) \rangle_n \right] f^3 \text{ a.u.} \quad ,$$

and the total correction $\delta E_n^\infty = \delta_{1p} E_n^\infty + \delta_{2p} E_n^\infty + \delta_s^{ee} E_n|_{\alpha^3} + \langle H_{vp} \rangle_n$ is

$$\delta E_n^\infty = \alpha^3 \left[\frac{4}{3} \left(\frac{19}{30} - \ln f \alpha^2 - \beta_n \right) \langle \delta^3(r_1) + \delta^3(r_2) \rangle_n + \left(\frac{14}{3} \ln f \alpha + \frac{164}{15} \right) \langle \delta^3(r) \rangle_n - \frac{7}{6\pi} Q_{12}^n \right] f^3 \text{ a.u. .}$$

Table captions

Table I. Elementary operators $\{E_k, k = 1, 15\}$ used in the calculation of the expectation values. n_k denotes the number of distinct polynomial terms generated by the action of E_k on a basis state.

Table II. The ground state expectation values $E_g = \langle H_0 \rangle$, $\langle H_0^2 \rangle$ and $\sigma^2 = \langle H_0^2 \rangle - E_g^2$ as a function of the basis dimension N_b .

Table III. Ground state expectation values of the singular differential operators $\langle \nabla_1^4 \rangle_E$, $\langle \nabla_1^4 \rangle$ and $\langle \nabla_3^4 \rangle_E$ as a function of the basis dimension N_b .

Table IV. Ground state expectation values of the singular Dirac distributions $\langle \delta(\vec{R}_1) \rangle$ and $\langle \delta(\vec{R}) \rangle$ as a function of the basis dimension N_b .

Table V. The ground state expectation values Q_{12}^g and Q_{13}^g defined by $Q_{ij}^g = \lim_{a \rightarrow 0} \langle \theta(r_{ij} - a) / r_{ij}^3 + 4\pi[\gamma + \ln(a/a_\mu)] \delta^3(r_{ij}) \rangle$, as a function of the basis dimension N_b .

Table I

k	n_k	E_k
1	4	$\hat{r}_1 \cdot \hat{r}_2 \partial_{r_1 r_2}^2$
2	4	$r^{-1}(r_2 \hat{r}_1 \cdot \hat{r}_2 - r_1) \partial_{r_1 r}^2$
3	4	$r^{-1}(r_1 \hat{r}_1 \cdot \hat{r}_2 - r_2) \partial_{r_2 r}^2$
4	1	$\partial_r^2 + 2r^{-1} \partial_r$
5	1	$r_1(r_2 r)^{-1} \hat{r}_1 \cdot \nabla_2^Y \partial_r$
6	1	$r_2(r_1 r)^{-1} \hat{r}_2 \cdot \nabla_1^Y \partial_r$
7	4	$r_2^{-1} \hat{r}_1 \cdot \nabla_2^Y \partial_{r_1} + r_1^{-1} \hat{r}_2 \cdot \nabla_1^Y \partial_{r_2}$
8	1	$(r_1 r_2)^{-1} \nabla_1^Y \cdot \nabla_2^Y$
9	3	$\partial_{r_1}^2 + 2r_1^{-1} \partial_{r_1}$
10	3	$\partial_{r_2}^2 + 2r_2^{-1} \partial_{r_2}$
11	1	$r_1^{-2} \nabla_1^Y \cdot \nabla_1^Y$
12	1	$r_2^{-2} \nabla_2^Y \cdot \nabla_2^Y$
13	3	$r_1^{-1} + r_2^{-1} - r^{-1}$
14	1	$-l_1(l_1 + 1)r_1^{-2}$
15	1	$-l_2(l_2 + 1)r_2^{-2}$

Table II

N_b	$E_g[\text{Ry}]$	$\langle H_0^2 \rangle [\text{Ry}^2]$	$\sigma^2 \times 10^{14} [\text{Ry}^2]$
324	−0.524 010 140 413 399 000 28	0.274 586 632 449 596	519 352.588 780 183 9
411	−0.524 010 140 455 551 566 88	0.274 586 628 565 868	126 562.113 711 645 6
512	−0.524 010 140 464 139 040 54	0.274 586 627 626 769	31 752.227 055 472 07
630	−0.524 010 140 465 665 621 87	0.274 586 627 375 932	6 508.536 436 039 192
764	−0.524 010 140 465 918 375 12	0.274 586 627 323 102	1 199.051 083 460 867
918	−0.524 010 140 465 954 391 13	0.274 586 627 313 704	255.472 732 778 647 0
1089	−0.524 010 140 465 959 038 66	0.274 586 627 311 885	73.086 762 209 023 06
1283	−0.524 010 140 465 960 002 45	0.274 586 627 311 421	26.589 554 852 763 07
1495	−0.524 010 140 465 960 160 85	0.274 586 627 311 222	6.672 153 792 304 994
1733	−0.524 010 140 465 960 203 19	0.274 586 627 311 175	1.965 916 264 831 472
1990	−0.524 010 140 465 960 212 96	0.274 586 627 311 165	0.965 692 558 621 058
2276	−0.524 010 140 465 960 214 82	0.274 586 627 311 160	0.465 897 417 244 748
2528	−0.524 010 140 465 960 215 25	0.274 586 627 311 158	0.308 152 352 372 668
<i>eff</i>	−0.524 010 140 465 960 215 4(6)	0.274 586 627 311 156(4)	0.23(23)

Table III

N_b	$\langle \nabla_1^4 \rangle_E$	$\langle \nabla_1^4 \rangle$	$\langle \nabla_3^4 \rangle_E$
324	2.532 451 004 442 6	2.532 445 719 29	5.255 396 862 891
411	2.532 451 050 420 6	2.532 451 697 56	5.255 397 122 254
512	2.532 451 056 877 0	2.532 450 741 84	5.255 397 117 353
630	2.532 451 009 132 0	2.532 449 964 61	5.255 397 051 034
764	2.532 451 018 719 1	2.532 450 992 21	5.255 397 086 467
918	2.532 451 022 453 6	2.532 451 056 52	5.255 397 094 127
1089	2.532 451 021 529 7	2.532 451 023 43	5.255 397 091 672
1283	2.532 451 020 589 3	2.532 451 019 49	5.255 397 091 024
1495	2.532 451 020 595 0	2.532 451 022 17	5.255 397 090 993
1733	2.532 451 020 587 2	2.532 451 020 24	5.255 397 090 958
1990	2.532 451 020 559 2	2.532 451 019 92	5.255 397 090 940
2276	2.532 451 020 559 6	2.532 451 020 43	5.255 397 090 949
2528	2.532 451 020 560 0	2.532 451 020 42	5.255 397 090 945
<i>eff</i>	2.532 451 020 559 6(3)	2.532 451 020 2(2)	5.255 397 090 945(4)

Table IV

N_b	$\langle\delta(\vec{R}_1)\rangle[a_0^{-3}]$	$\langle\delta(\vec{R})\rangle[a_0^{-3}]$
324	0.020 733 174 230 2	0.000 171 000 000 8
411	0.020 733 203 838 1	0.000 170 999 383 2
512	0.020 733 199 804 5	0.000 170 999 967 2
630	0.020 733 193 292 2	0.000 170 997 306 7
764	0.020 733 197 986 7	0.000 170 996 885 4
918	0.020 733 198 238 9	0.000 170 996 811 0
1089	0.020 733 198 094 3	0.000 170 996 832 4
1283	0.020 733 197 999 5	0.000 170 996 756 0
1495	0.020 733 198 024 3	0.000 170 996 767 3
1733	0.020 733 198 007 4	0.000 170 996 760 1
1990	0.020 733 198 003 4	0.000 170 996 757 7
2276	0.020 733 198 005 3	0.000 170 996 757 1
2528	0.020 733 198 005 0	0.000 170 996 756 8
<i>eff</i>	0.020 733 198 004 6(8)	0.000 170 996 756 7(4)

Table V

N_b	$Q_{12}^g[a_\mu^{-3}]$	$Q_{13}^g[a_\mu^{-3}]$
324	0.095 757 780 75	-2.776 563 295
411	0.095 757 975 79	-2.776 588 343
512	0.095 757 804 27	-2.776 583 829
630	0.095 758 749 78	-2.776 578 687
764	0.095 758 904 03	-2.776 582 810
918	0.095 758 930 40	-2.776 582 894
1089	0.095 758 918 40	-2.776 582 776
1283	0.095 758 949 78	-2.776 582 694
1495	0.095 758 944 76	-2.776 582 722
1733	0.095 758 947 86	-2.776 582 703
1990	0.095 758 949 04	-2.776 582 700
2276	0.095 758 949 31	-2.776 582 702
<i>eff</i>	0.095 758 949 4(3)	-2.776 582 702(1)

References

- [1] E. A. Hylleraas, Z. Phys. **48** 469 (1928); **54** 347 (1929).
- [2] G. W. F. Drake, Phys. Rev. A **18** 820 (1978).
- [3] G. W. F. Drake, *High Precision Calculations for the Rydberg States of Helium*, in Long-Range Casimir Forces: Theory and Recent Experiments on Atomic Systems, Ed. by F. S. Levin and D. A. Micha. Plenum Press, New York, 1993, p. 107.
- [4] G. W. F. Drake, *High Precision Calculations for Helium*, in Atomic, Molecular and Optical Physics Handbook, Ed. by G. W. F. Drake. AIP Press, Woodbury, New York, 1996, p. 154.
- [5] P. A. M. Dirac, Canad. J. Math. **2** 129 (1950).
- [6] D. G. Currie, T. F. Jordan and E. C. G. Sudarshan, Rev. Mod. Phys. **35** 350 (1962).
- [7] R. N. Hill, *The Origins of Predictive Relativistic Mechanics*, in Relativistic Action at a Distance: Classical and Quantum Aspects, Ed. by J. Llosa, Springer-Verlag, Berlin (1982), p. 104.
- [8] G. N. Fleming, Phys. Rev. **137** B188 (1965).
- [9] H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One - and Two Electron Systems*, Plenum Publishing Corporation, New York (1977).
- [10] P. Strange, *Relativistic Quantum Mechanics* , Cambridge University Press (1998), p. 322.
- [11] J. A. Wheeler, Ann. N. Y. Acad. Sci. **48**, 219 (1946).
- [12] A. P. Mills Jr., Phys. Rev. Lett. **46** 717 (1981); **50** 671 (1983).
- [13] G. Ferrante, Phys. Rev. **170**, 76 (1968).
- [14] Y. K. Ho, Phys. Lett. A **144**, 237 (1990).
- [15] A. M. Frolov, Phys. Rev. A **60** 2834 (1999).

- [16] R. Krivec, V. B. Mandelzweig and K. Varga, Phys. Rev. **A**, 062503-1 (2000).
- [17] G. W. F. Drake, M. M. Casar and R. A. Nistor, Phys. Rev. A **65** 054501-1 (2002).
- [18] Y. K. Ho, Phys. Rev. A **19**, 2347 (1979).
- [19] J. Usukura and Y. Suzuki, Phys. Rev. A **66** 010502-1 (2002).
- [20] J. Sapirstein, *Quantum Electrodynamics*, in Atomic, Molecular and Optical Physics Handbook, Ed. by G. W. F. Drake. AIP Press, Woodbury, New York, 1996, p. 327.
- [21] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics*, McGraw-Hill Book Company, New York (1964).
- [22] W. Heisenberg, Zeits. f. Physik **90** 209 (1934).
- [23] E. A. Uehling, Phys. Rev. **48** 55 (1935).
- [24] W. E. Lamb Jr. and R. C. Retherford, Phys. Rev. **72**, 241 (1947).
- [25] T. Fulton and P. C. Martin, Phys. Rev. **95**, 811 (1954).
- [26] H. Araki, Prog. Theor. Phys. **17**, 619 (1957).
- [27] G. W. F. Drake and S. P. Goldman, Can. J. Phys. **77** 835 (1999).
- [28] K. Pachucki, J. Phys. B: At. Mol. Opt. Phys. **31**, 5123 (1998).
- [29] R. Foot and S. N. Gninenko, Phys. Lett. B **480** 171 (2000).
- [30] R. Karplus and A. Klein, Phys. Rev. **87** 848 (1952).
- [31] E. D. Therot Jr., R. H. Beers, V. H. Hughes and K. O. H. Ziock, Phys. Rev. A **2**, 707 (1970).
- [32] A. K. Bhatia and R. J. Drachman, Phys. Rev. A **28** 2523 (1983).